



LUND
UNIVERSITY



$\pi\pi$ and πK at Two Loops

Johan Bijnens

Lund University

bijnens@thep.lu.se

<http://www.thep.lu.se/~bijnens/chpt.html>

Overview

- $\pi\pi$ Scattering in Three Flavour ChPT, J. Bijnens, P. Dhonte and P. Talavera, hep-ph/0401039, JHEP 0401(2004)050
- πK Scattering in Three Flavour ChPT, J. Bijnens, P. Dhonte and P. Talavera hep-ph/0404150, JHEP 0405(2004)036
- Introduction
- Why (Effective) Field Theory
- Chiral Perturbation Theory
- Two Loop: General
- Two Loop: Three Flavours
 - General fitting strategy and some comments
 - $\pi\pi, \pi K$
- Conclusions

Introduction

- $\pi\pi$ and πK scattering are basic strong processes
Study them as precise as possible
- Earlier: $\pi\pi$ proved that two-flavour case of $\langle \bar{q}q \rangle$
- Three flavour case: works or problems with strange quark loops (in scalar sector)?
- πK excellent place to study this
- Need precise calculations also here

Why (Effective) Field Theory?

- Effective:**
- Use right degrees of freedom : essence of (most) physics
 - Gap in the spectrum \implies separation of scales
 - With lower d.o.f.: build most general Lagrangian

⇒ ∞ parameters
⇒ Where did predictivity go ? } \implies power counting

Why (Effective) Field Theory?

Field Theory

- ➡ Only known way to combine QM and special relativity
- ➡ Taylor series does not work (convergence radius zero)
- ➡ Continuum of excitation states to be taken into account
- ➡ Off-shell effects fully under control: these effects are there as new free parameters
- ➡ model-independent and systematic: ALL effects at given order included
- ➡ Theory \implies errors can be estimated
- ➡ Many parameters (but possible modelspace is large)
- ➡ Expansion might not converge (often still useful for model classification)

Chiral Perturbation Theory

Degrees of freedom: Goldstone Bosons from Chiral Symmetry Spontaneous Breakdown

Power counting: Dimensional counting


Expected breakdown scale: Resonances, so M_ρ or higher depending on the channel

Chiral Symmetry

QCD: 3 light quarks: equal mass: interchange: $SU(3)_V$

But
$$\mathcal{L}_{QCD} = \sum_{q=u,d,s} [i\bar{q}_L \not{D} q_L + i\bar{q}_R \not{D} q_R - m_q (\bar{q}_R q_L + \bar{q}_L q_R)]$$

So if $m_q = 0$ then $SU(3)_L \times SU(3)_R$.

Can also see that via  $v < c, m_q \neq 0 \implies$
 $v = c, m_q = 0 \not\implies$


Chiral Perturbation Theory

$$\langle \bar{q}q \rangle = \langle \bar{q}_L q_R + \bar{q}_R q_L \rangle \neq 0$$


$SU(3)_L \times SU(3)_R$ broken spontaneously to $SU(3)_V$

8 generators broken \implies 8 massless degrees of freedom **and**
interaction vanishes at zero momentum

Power counting in momenta:




$$p^2$$



$$(p^2)^2 (1/p^2)^2 p^4 = p^4$$



$$1/p^2$$



$$(p^2) (1/p^2) p^4 = p^4$$

$$\int d^4 p$$

$$p^4$$

Two Loop: General

Lagrangian Structure:

	2 flavour		3 flavour		3+3 PQChPT	
p^2	F, B	2	F_0, B_0	2	F_0, B_0	2
p^4	l_i^r, h_i^r	7+3	L_i^r, H_i^r	10+2	\hat{L}_i^r, \hat{H}_i^r	11+2
p^6	c_i^r	53+4	C_i^r	90+4	K_i^r	112+3

p^2 : Weinberg 1966

p^4 : Gasser, Leutwyler 84,85

p^6 : JB, Colangelo, Ecker 99,00

Note {

- ➡ PQ \implies Talk by Timo Lähde
- ➡ All infinities known
- ➡ Two Flavour Most Things Done

Three Flavours at Two Loop

$\Pi_{VV\pi}, \Pi_{VV\eta}, \Pi_{VVK}$	Kambor, Golowich; Kambor, Dürr; Amorós, JB, Talavera
$\Pi_{VV\rho\omega}$	Maltman
$\Pi_{AA\pi}, \Pi_{AA\eta}, F_\pi, F_\eta, m_\pi, m_\eta$	Kambor, Golowich; Amorós, JB, Talavera
Π_{SS}	Moussallam L_4^r, L_6^r
$\Pi_{VVK}, \Pi_{AAK}, F_K, m_K$	Amorós, JB, Talavera
$K_{\ell 4}, \langle \bar{q}q \rangle$	Amorós, JB, Talavera L_1^r, L_2^r, L_3^r
$F_M, m_M, \langle \bar{q}q \rangle (m_u \neq m_d)$	Amorós, JB, Talavera $L_{5,7,8}^r, m_u/m_d$
$F_{V\pi}, F_{VK^+}, F_{VK^0}$	Post, Schilcher; JB, Talavera L_9^r
$K_{\ell 3}$	Post, Schilcher; JB, Talavera V_{us}
$F_{S\pi}, F_{SK}$ (includes σ -terms)	JB, Dhonte L_4^r, L_6^r
$K, \pi \rightarrow \ell\nu\gamma$	Geng, Ho, Wu L_{10}^r
$\pi\pi$	JB, Dhonte, Talavera
πK	JB, Dhonte, Talavera

General Strategy and some comments

- Find enough inputs from experiment
- C_i^r :
 - kinematical dependence: agree well with single resonance saturation
 - quark mass+kinematical: if vector dominated, seems to be OK
 - quark mass+kinematical: if scalar dominated: which scalars? (not σ)
 - quark masses: which scalars? unrealistically large estimates
- in p^6 physical or lowest order masses: thresholds in right place requires physical

General Strategy and some comments

Inputs:

$$K_{\ell 4}: F(0), G(0), \lambda$$

$$m_{\pi^0}^2, m_{\eta}^2, m_{K^+}^2, m_{K^0}^2$$

$$F_{\pi^+}$$

$$F_{K^+}/F_{\pi^+}$$

E865 BNL
em with Dashen violation

General Strategy and some comments

Inputs:

$$K_{\ell 4}: F(0), G(0), \lambda$$

$$m_{\pi^0}^2, m_{\eta}^2, m_{K^+}^2, m_{K^0}^2$$

$$F_{\pi^+}$$

$$F_{K^+}/F_{\pi^+}$$

$$m_s/\hat{m}$$

$$L_4^r, L_6^r$$

24 (26)

$$\hat{m} = (m_u + m_d)/2$$

Vary \Rightarrow other L_i^r vary correlated

E865 BNL
em with Dashen violation

General Strategy and some comments

Inputs:

$$K_{\ell 4}: F(0), G(0), \lambda$$

$$m_{\pi^0}^2, m_{\eta}^2, m_{K^+}^2, m_{K^0}^2$$

$$F_{\pi^+}$$

$$F_{K^+}/F_{\pi^+}$$

$$m_s/\hat{m}$$


$$24 \quad (26)$$

$$\hat{m} = (m_u + m_d)/2$$


$$L_4^r, L_6^r$$

Vary \Rightarrow other L_i^r vary correlated

C_i^r from single resonance approximation



$$|q^2| \ll m_{\rho}^2, m_S^2$$



General Strategy: fit results



	fit 10	same p^4	fit B	fit D
$10^3 L_1^r$	0.43 ± 0.12	0.38	0.44	0.44
$10^3 L_2^r$	0.73 ± 0.12	1.59	0.60	0.69
$10^3 L_3^r$	-2.53 ± 0.37	-2.91	-2.31	-2.33
$10^3 L_4^r$	$\equiv 0$	$\equiv 0$	$\equiv 0.5$	$\equiv 0.2$
$10^3 L_5^r$	0.97 ± 0.11	1.46	0.82	0.88
$10^3 L_6^r$	$\equiv 0$	$\equiv 0$	$\equiv 0.1$	$\equiv 0$
$10^3 L_7^r$	-0.31 ± 0.14	-0.49	-0.26	-0.28
$10^3 L_8^r$	0.60 ± 0.18	1.00	0.50	0.54

- errors are very correlated
- $\mu = 770$ MeV; 550 or 1000 within errors
- varying C_i^r factor 2 about errors
- $L_4^r, L_6^r \approx -0.3, \dots, 0.6 \cdot 10^{-3}$ OK
- fit B**: small corrections to pion “sigma” term, fit scalar radius
- fit D**: fit $\pi\pi$ and πK thresholds

General Strategy: some outputs

	fit 10	same p^4	fit B	fit D
$2B_0\hat{m}/m_\pi^2$	0.736	0.991	1.129	0.958
$m_\pi^2: p^4, p^6$	0.006,0.258	0.009, $\equiv 0$	-0.138,0.009	-0.091,0.133
$m_K^2: p^4, p^6$	0.007,0.306	0.075, $\equiv 0$	-0.149,0.094	-0.096,0.201
$m_\eta^2: p^4, p^6$	-0.052,0.318	0.013, $\equiv 0$	-0.197,0.073	-0.151,0.197
m_u/m_d	0.45 ± 0.05	0.52	0.52	0.50
F_0 [MeV]	87.7	81.1	70.4	80.4
$F_K/F_\pi: p^4, p^6$	0.169,0.051	0.22, $\equiv 0$	0.153,0.067	0.159,0.061

- ➡ Pattern of mass corrections can vary a lot
- ➡ F_K/F_π always OK expansion
- ➡ $m_u = 0$ always very far from the fits
- ➡ F_0 : pion decay constant in the chiral limit





$$a_0^0 = 0.220 \pm 0.005, \quad a_0^2 = -0.0444 \pm 0.0010$$

Colangelo, Gasser, Leutwyler

$$a_0^0 = 0.159 \quad a_0^2 = -0.0454 \quad \text{at order } p^2$$



$\pi\pi$ subthreshold parameters



$$C_1 = 1.104 \pm 0.009, C_2 = 1.120 \pm 0.027$$

Colangelo, Gasser, Leutwyler

$$C_1 = C_2 = 1 \text{ at order } p^2$$





$$a_0^{1/2} = 0.224 \pm 0.022, \quad a_{3/2}^2 = -0.0448 \pm 0.0077$$

Büttiker, Descotes-Genon, Moussallam

$$a_0^{1/2} = 0.142 \quad a_0^2 = -0.0708 \quad \text{at order } p^2$$

πK subthreshold parameters



$$c_{10}^+ = 0.87 \pm 0.08, \quad c_{00}^- = 8.92 \pm 0.38$$

Büttiker, Descotes-Genon, Moussallam

πK subthreshold parameters

	Vector	Scalar	Sum Reso	chiral order	p^2	p^4	p^6
c_{00}^+	-0.02	0.13	0.11	2	0	0.122	0.007
c_{10}^+	0.018	-0.063	-0.045	2	0.5704	-0.113	0.460
c_{00}^-	0.21	0.17	0.38	2	8.070	0.311	0.017
c_{20}^+	-0.0053	0.0023	-0.0030	4	—	0.0256	-0.0254
c_{10}^-	-0.11	-0.04	-0.15	4	—	-0.0254	0.121
c_{01}^+	-0.27	0.28	0.01	4	—	1.667	1.492
c_{30}^+	0.00026	0.00010	0.00036	6	—	0.00121	0.00071
c_{20}^-	0.0037	0.00060	0.0043	6	—	0.00478	0.00320
c_{11}^+	0.017	-0.008	0.009	6	—	-0.126	-0.006
c_{01}^-	0.25	0.04	0.29	6	—	0.229	0.196


Resonance contributions, units: $m_{\pi^+}^{2i+2j}$ (c_{ij}^+) and $m_{\pi^+}^{2i+2j+1}$ (c_{ij}^-)

Chiral order at which they first have tree level contributions

Contributions with the $L_i^r = C_i^r = 0$ at $\mu = 0.77$ GeV.


πK subthreshold parameters

	Fit 10	BDM	Lang
c_{00}^+	0.278	2.01 ± 1.10	-0.52 ± 2.03
c_{10}^+	0.898	0.87 ± 0.08	0.55 ± 0.07
c_{00}^-	8.99	8.92 ± 0.38	7.31 ± 0.90
c_{20}^+	0.003	0.024 ± 0.006	
c_{10}^-	0.088	0.31 ± 0.01	0.21 ± 0.04
c_{01}^+	3.8	2.07 ± 0.10	2.06 ± 0.22
c_{30}^+	0.0025	0.0034 ± 0.0008	
c_{20}^-	0.013	0.0085 ± 0.0001	
c_{11}^+	-0.10	-0.066 ± 0.010	
c_{01}^-	0.71	0.62 ± 0.06	0.51 ± 0.10
c_{02}^+	0.23	0.34 ± 0.03	




$\pi\pi$ and πK

$\pi\pi$ constraints




$\pi\pi$ and πK

$\pi\pi$ constraints




πK constraints




$\pi\pi$ and πK

$\pi\pi$ constraints



πK constraints



preferred region: fit D: $10^3 L_4^r \approx 0.2$, $10^3 L_6^r \approx 0.0$

Conclusions

Three flavour ChPT at 2 loops doing fine: much progress

- many calculations done
- things seem to work but convergence is fairly slow
- “kinematical” and “vector” C_i^r seem to be OK
- L_4^r, L_6^r nonzero but reasonable for large N_c
- $\eta \rightarrow 3\pi$, isobreaking in $K_{\ell 3}$: parts done

Conclusions

Three flavour ChPT at 2 loops doing fine: much progress

- many calculations done
- things seem to work but convergence is fairly slow
- “kinematical” and “vector” C_i^r seem to be OK
- L_4^r, L_6^r nonzero but reasonable for large N_c
- $\eta \rightarrow 3\pi$, isobreaking in $K_{\ell 3}$: parts done

πK open problems

- Cleaning up C_i^r contributions and uncertainties
- Properly predicting threshold parameters