



LUND
UNIVERSITY



PARTIALLY QUENCHED AND THREE FLAVOURS AT TWO LOOPS

Johan Bijnens

Lund University

`bijnens@thep.lu.se`

`http://www.thep.lu.se/~bijnens`

Various ChPT: `http://www.thep.lu.se/~bijnens/chpt.html`

Overview

- Chiral Perturbation Theory
- Lagrangians
- What is partially quenched?
- PQChPT and some problems
- Some results of PQChPT at Two-loops
- A Two-loop Timeline
- Some comments on and results from ChPT at two loops
- Conclusions

Chiral Perturbation Theory

Degrees of freedom: Goldstone Bosons from Chiral Symmetry Spontaneous Breakdown

Power counting: Dimensional counting

Expected breakdown scale: Resonances, so M_ρ or higher depending on the channel

Chiral Symmetry

QCD: 3 light quarks: equal mass: interchange: $SU(3)_V$

But
$$\mathcal{L}_{QCD} = \sum_{q=u,d,s} [i\bar{q}_L \not{D} q_L + i\bar{q}_R \not{D} q_R - m_q (\bar{q}_R q_L + \bar{q}_L q_R)]$$

So if $m_q = 0$ then $SU(3)_L \times SU(3)_R$.


Chiral Perturbation Theory

$$\langle \bar{q}q \rangle = \langle \bar{q}_L q_R + \bar{q}_R q_L \rangle \neq 0$$


$SU(3)_L \times SU(3)_R$ broken spontaneously to $SU(3)_V$

8 generators broken \implies 8 massless degrees of freedom
and interaction vanishes at zero momentum

Power counting in momenta:




$$p^2$$



$$(p^2)^2 (1/p^2)^2 p^4 = p^4$$



$$1/p^2$$



$$(p^2) (1/p^2) p^4 = p^4$$

$$\int d^4 p$$

$$p^4$$

Two Loop: Lagrangians

Lagrangian Structure:

	2 flavour		3 flavour		3+3 PQChPT	
p^2	F, B	2	F_0, B_0	2	F_0, B_0	2
p^4	l_i^r, h_i^r	7+3	L_i^r, H_i^r	10+2	\hat{L}_i^r, \hat{H}_i^r	11+2
p^6	c_i^r	53+4	C_i^r	90+4	K_i^r	112+3

p^2 : Weinberg 1966

p^4 : Gasser, Leutwyler 84,85

p^6 : JB, Colangelo, Ecker 99,00

Eurodaphne II/Euridice (PQChPT)

Constructing Lagrangians

Easy part:

- Construct a complete set of needed basic objects
- Often useful to make everything transform under H alone (conserved part)
- Construct all terms up to the order you want

Tricky part: finding a minimal basis

- Field redefinitions or equations of motion
- Partial integration
- Cayley-Hamilton identities
- Bianchi identities, $\det \chi$ is also chirally invariant

Final check: can I determine all from experiment?

Constructing Lagrangians

Infinites:

the general divergence structure can be derived using heat kernel methods and/or background field methods.

Very useful for checks on calculations

What is Partially Quenched?

In Lattice gauge theory one calculates

$$\langle 0 | (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) | \rangle$$

$$= \frac{\int [dq][d\bar{q}][dG] (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}}}{\int [dq][d\bar{q}][dG] \bar{d}\gamma_5 u(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}}}$$

for Euclidean separations x

Integrals also performed after rotation to Euclidean
(note that I use Minkowski notation throughout)

What is Partially Quenched?

$$\int [dq][d\bar{q}][dG] (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}} \propto$$

$$\int [dG] e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}} (\not{D}_G^u)^{-1}(x, 0) (\not{D}_G^d)^{-1}(0, x) \det(\not{D}_G)_{\text{QCD}}$$

What is Partially Quenched?

$$\int [dq][d\bar{q}][dG] (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}} \propto$$

$$\int [dG] e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}} (\not{D}_G^u)^{-1}(x, 0) (\not{D}_G^d)^{-1}(0, x) \det(\not{D}_G)_{\text{QCD}}$$

$\int [dG]$ done via importance sampling

What is Partially Quenched?

$$\int [dq][d\bar{q}][dG] (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}} \propto$$

$$\int [dG] e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}} (\not{D}_G^u)^{-1}(x, 0) (\not{D}_G^d)^{-1}(0, x) \det(\not{D}_G)_{\text{QCD}}$$

$\int [dG]$ done via importance sampling

- Quenched: get distribution from $e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}}$ only

What is Partially Quenched?

$$\int [dq][d\bar{q}][dG] (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}} \propto$$

$$\int [dG] e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}} (\not{D}_G^u)^{-1}(x, 0) (\not{D}_G^d)^{-1}(0, x) \det(\not{D}_G)_{\text{QCD}}$$

$\int [dG]$ done via importance sampling

- Quenched: get distribution from $e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}}$ only
- Unquenched: include $\det(\not{D}_G)_{\text{QCD}}$ VERY expensive

What is Partially Quenched?

$$\int [dq][d\bar{q}][dG] (\bar{u}\gamma_5 d)(x) (\bar{d}\gamma_5 u)(0) e^{i \int d^4 y \mathcal{L}_{\text{QCD}}} \propto$$

$$\int [dG] e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}} (\not{D}_G^u)^{-1}(x, 0) (\not{D}_G^d)^{-1}(0, x) \det(\not{D}_G)_{\text{QCD}}$$

$\int [dG]$ done via importance sampling

- Quenched: get distribution from $e^{i \int d^4 x (-1/4) G_{\mu\nu} G^{\mu\nu}}$ only
- Unquenched: include $\det(\not{D}_G)_{\text{QCD}}$ VERY expensive
- Partially quenched: $(\not{D}_G^u)^{-1}(x, 0) (\not{D}_G^d)^{-1}(0, x)$
DIFFERENT Quarks then in $\det(\not{D}_G)_{\text{QCD}}$

What is Partially Quenched?

Why do this?

- Is not Quenched: Real QCD is continuous limit from Partially Quenched
- More handles to turn:
 - Allows more systematic studies by varying parameters
 - Sometimes allows to disentangle things from different observables
- $\det(\not{D}_G)_{\text{QCD}}$: Sea quarks
- $(\not{D}_G^u)^{-1}(x, 0)(\not{D}_G^d)^{-1}(0, x)$: Valence Quarks

What is Partially Quenched?

Why not do this?


- It is not QCD as soon as $\text{Valence} \neq \text{Sea}$
- not a Quantum Field Theory
- No unitarity
- No clusterdecomposition
- No CPT theorem
- No spin statistics theorem

What is Partially Quenched?

Do anyway

- Cheap computationally
- Allows extra studies
- Hope it works near real QCD case
- Is a well defined Euclidean statistical model

ChPT and Lattice QCD



Valence is *easy* to deal with in lattice QCD


Sea is *very difficult*

They can be treated separately: i.e. different quark masses

Partially Quenched ChPT (PQChPT)

PQChPT at Two Loops: General

Add ghost quarks: remove the unwanted free valence loops



Possible problem: QCD \implies ChPT relies heavily on unitarity

Partially quenched: at least one dynamical sea quark
 $\implies \Phi_0$ is heavy: remove from PQChPT

Symmetry group becomes $SU(n_v + n_s | n_v) \times SU(n_v + n_s | n_v)$
 (approximately)

PQChPT at Two Loops: General

Essentially all manipulations from ChPT go through to PQChPT when changing trace to supertrace and adding fermionic variables

Exceptions: baryons and Cayley-Hamilton relations

So Luckily: can use the n flavour work in ChPT at two loop order to obtain for PQChPT: Lagrangians and infinities

Very important note: ChPT is a limit of PQChPT
 \implies LECs from ChPT are linear combinations of LECs of PQChPT with the **same** number of sea quarks.

$$\text{E.g. } L_1^r = L_0^{r(3pq)} / 2 + L_1^{r(3pq)}$$

PQChPT at Two Loop

Subject started:

valence equal mass, 3 sea equal mass:

$m_{\pi^+}^2$: JB, Danielsson, Lähde, hep-lat/0406017

Other mass combinations:

F_{π^+} : JB, Lähde, hep-lat/0501014

F_{π^+} , $m_{\pi^+}^2$ **two sea quarks**: JB, Lähde, hep-lat/0506004

$m_{\pi^+}^2$: JB, Danielsson, Lähde, hep-lat/0602003







Neutral masses: JB, Danielsson, hep-lat/0606017

Actual Calculations: {


- heavy use of FORM **Vermaseren**
- use PQ without super Φ_0 in super-symmetric formalism
- Main problem: sheer size of the expressions

Iso breaking from lattice data: a and L extrapolations needed

Long Expressions



Long Expressions



$$\begin{aligned}
 \delta_{\text{loops}}^{(6)22} = & \pi_{16} L_0^2 [4/9 \chi_\eta \chi_4 - 1/2 \chi_1 \chi_3 + \chi_{13}^2 - 13/3 \bar{\chi}_1 \chi_{13} - 35/18 \bar{\chi}_2] - 2 \pi_{16} L_1^2 \chi_{13}^2 \\
 & - \pi_{16} L_2^2 [11/3 \chi_\eta \chi_4 + \chi_{13}^2 + 13/3 \bar{\chi}_2] + \pi_{16} L_3^2 [4/9 \chi_\eta \chi_4 - 7/12 \chi_1 \chi_3 + 11/6 \chi_{13}^2 - 17/6 \bar{\chi}_1 \chi_{13} - 43/36 \bar{\chi}_2] \\
 & + \pi_{16}^2 [-15/64 \chi_\eta \chi_4 - 59/384 \chi_1 \chi_3 + 65/384 \chi_{13}^2 - 1/2 \bar{\chi}_1 \chi_{13} - 43/128 \bar{\chi}_2] - 48 L_4^2 L_5^2 \bar{\chi}_1 \chi_{13} - 72 L_4^2 \bar{\chi}_2^2 \\
 & - 8 L_5^2 \chi_{13}^2 + \bar{A}(\chi_p) \pi_{16} [-1/24 \chi_p + 1/48 \bar{\chi}_1 - 1/8 \bar{\chi}_1 R_{\eta\eta}^c + 1/16 \bar{\chi}_1 R_p^c - 1/48 R_{\eta\eta}^c \chi_p - 1/16 R_{\eta\eta}^c \chi_q \\
 & + 1/48 R_{pp}^c \chi_\eta + 1/16 R_p^c \chi_{13}] + \bar{A}(\chi_p) L_0^2 [8/3 R_{\eta\eta}^c \chi_p + 2/3 R_p^c \chi_p + 2/3 R_{\eta\eta}^c] + \bar{A}(\chi_p) L_5^2 [2/3 R_{\eta\eta}^c \chi_p \\
 & + 5/3 R_p^c \chi_p + 5/3 R_{\eta\eta}^c] + \bar{A}(\chi_p) L_4^2 [-2 \bar{\chi}_1 \bar{\chi}_{\eta\eta}^{pp} - 2 \bar{\chi}_1 R_{\eta\eta}^c + 3 \bar{\chi}_1 R_p^c] + \bar{A}(\chi_p) L_5^2 [-2/3 \bar{\chi}_{\eta\eta}^{pp} - R_{\eta\eta}^c \chi_p \\
 & + 1/3 R_{\eta\eta}^c \chi_q + 1/2 R_p^c \chi_p - 1/6 R_p^c \chi_q] + \bar{A}(\chi_p)^2 [1/16 + 1/72 (R_{\eta\eta}^c)^2 - 1/72 R_{\eta\eta}^c R_p^c + 1/288 (R_p^c)^2] \\
 & + \bar{A}(\chi_p) \bar{A}(\chi_{ps}) [-1/36 R_{\eta\eta}^c - 5/72 R_{\eta\eta}^c + 7/144 R_p^c] - \bar{A}(\chi_p) \bar{A}(\chi_{qs}) [1/36 R_{\eta\eta}^c + 1/24 R_{\eta\eta}^c + 1/48 R_p^c] \\
 & + \bar{A}(\chi_p) \bar{A}(\chi_\eta) [-1/72 R_{\eta\eta}^c R_{\eta\eta}^c + 1/144 R_p^c R_{\eta\eta}^c] + 1/8 \bar{A}(\chi_p) \bar{A}(\chi_{13}) + 1/12 \bar{A}(\chi_p) \bar{A}(\chi_{46}) R_{pp}^c \\
 & + \bar{A}(\chi_p) \bar{B}(\chi_p, \chi_p; 0) [1/4 \chi_p - 1/18 R_{\eta\eta}^c R_p^c \chi_p - 1/72 R_{\eta\eta}^c R_p^c + 1/18 (R_p^c)^2 \chi_p + 1/144 R_p^c R_{\eta\eta}^c] \\
 & + \bar{A}(\chi_p) \bar{B}(\chi_p, \chi_\eta; 0) [1/18 R_{\eta\eta}^c R_p^c \chi_p - 1/18 R_{\eta\eta}^c R_p^c \chi_p] + \bar{A}(\chi_p) \bar{B}(\chi_q, \chi_q; 0) [-1/72 R_{\eta\eta}^c R_p^c + 1/144 R_p^c R_{\eta\eta}^c] \\
 & - 1/12 \bar{A}(\chi_p) \bar{B}(\chi_{ps}, \chi_{ps}; 0) R_{\eta\eta}^c R_p^c \chi_p - 1/18 \bar{A}(\chi_p) \bar{B}(\chi_1, \chi_3; 0) R_{\eta\eta}^c R_p^c \chi_p \\
 & + 1/18 \bar{A}(\chi_p) \bar{C}(\chi_p, \chi_p, \chi_p; 0) R_{\eta\eta}^c R_p^c \chi_p + \bar{A}(\chi_p; \varepsilon) \pi_{16} [1/8 \bar{\chi}_1 R_{\eta\eta}^c - 1/16 \bar{\chi}_1 R_p^c - 1/16 R_{\eta\eta}^c \chi_p - 1/16 R_{\eta\eta}^c] \\
 & + \bar{A}(\chi_{ps}) \pi_{16} [1/16 \chi_{ps} - 3/16 \chi_{qs} - 3/16 \bar{\chi}_1] - 2 \bar{A}(\chi_{ps}) L_0^2 \chi_{ps} - 5 \bar{A}(\chi_{ps}) L_3^2 \chi_{ps} - 3 \bar{A}(\chi_{ps}) L_4^2 \bar{\chi}_1 \\
 & + \bar{A}(\chi_{ps}) L_5^2 \chi_{13} + \bar{A}(\chi_{ps}) \bar{A}(\chi_\eta) [7/144 R_{\eta\eta}^c - 5/72 R_{\eta\eta}^c - 1/48 R_{\eta\eta}^c + 5/72 R_{\eta\eta}^c - 1/36 R_{\eta\eta}^c] \\
 & + \bar{A}(\chi_{ps}) \bar{B}(\chi_p, \chi_p; 0) [1/24 R_{\eta\eta}^c \chi_p - 5/24 R_{\eta\eta}^c \chi_{ps}] + \bar{A}(\chi_{ps}) \bar{B}(\chi_p, \chi_\eta; 0) [-1/18 R_{\eta\eta}^c R_{\eta\eta}^c \chi_p \\
 & - 1/9 R_{\eta\eta}^c R_{\eta\eta}^c \chi_{ps}] - 1/48 \bar{A}(\chi_{ps}) \bar{B}(\chi_q, \chi_q; 0) R_{\eta\eta}^c + 1/18 \bar{A}(\chi_{ps}) \bar{B}(\chi_1, \chi_3; 0) R_{\eta\eta}^c \chi_s \\
 & + 1/9 \bar{A}(\chi_{ps}) \bar{B}(\chi_1, \chi_3; 0, k) R_{\eta\eta}^c + 3/16 \bar{A}(\chi_{ps}; \varepsilon) \pi_{16} [\chi_s + \bar{\chi}_1] - 1/8 \bar{A}(\chi_{p4})^2 - 1/8 \bar{A}(\chi_{p4}) \bar{A}(\chi_{p6}) \\
 & + 1/8 \bar{A}(\chi_{p4}) \bar{A}(\chi_{46}) - 1/32 \bar{A}(\chi_{p6})^2 + \bar{A}(\chi_\eta) \pi_{16} [1/16 \bar{\chi}_1 R_{\eta\eta}^c + 1/48 R_{\eta\eta}^c \chi_\eta + 1/16 R_{\eta\eta}^c \chi_{13}] \\
 & + \bar{A}(\chi_\eta) L_0^2 [4 R_{\eta\eta}^c \chi_\eta + 2/3 R_{\eta\eta}^c \chi_\eta] - 8 \bar{A}(\chi_\eta) L_1^2 \chi_\eta - 2 \bar{A}(\chi_\eta) L_2^2 \chi_\eta + \bar{A}(\chi_\eta) L_3^2 [4 R_{\eta\eta}^c \chi_\eta + 5/3 R_{\eta\eta}^c \chi_\eta] \\
 & + \bar{A}(\chi_\eta) L_4^2 [4 \chi_\eta + \bar{\chi}_1 R_{\eta\eta}^c] - \bar{A}(\chi_\eta) L_5^2 [1/6 R_{\eta\eta}^c \chi_q + R_{\eta\eta}^c \chi_{13} + 1/6 R_{\eta\eta}^c \chi_\eta] + 1/288 \bar{A}(\chi_\eta)^2 (R_{\eta\eta}^c)^2 \\
 & + 1/12 \bar{A}(\chi_\eta) \bar{A}(\chi_{46}) R_{\eta\eta}^c + \bar{A}(\chi_\eta) \bar{B}(\chi_p, \chi_p; 0) [-1/36 \bar{\chi}_{\eta\eta}^{pp} - 1/18 R_{\eta\eta}^c R_{\eta\eta}^c \chi_p + 1/18 R_{\eta\eta}^c R_p^c \chi_p \\
 & + 1/144 R_{\eta\eta}^c R_{\eta\eta}^c] + \bar{A}(\chi_\eta) \bar{B}(\chi_p, \chi_\eta; 0) [-1/18 \bar{\chi}_{\eta\eta}^{pp} + 1/18 \bar{\chi}_{\eta\eta}^{pp} + 1/18 (R_{\eta\eta}^c)^2 R_{\eta\eta}^c \chi_p] \\
 & - 1/12 \bar{A}(\chi_\eta) \bar{B}(\chi_{ps}, \chi_{ps}; 0) R_{\eta\eta}^c \chi_{ps} - \bar{A}(\chi_\eta) \bar{B}(\chi_\eta, \chi_\eta; 0) [1/216 R_{\eta\eta}^c \chi_4 + 1/27 R_{\eta\eta}^c \chi_6] \\
 & - 1/18 \bar{A}(\chi_\eta) \bar{B}(\chi_1, \chi_3; 0) R_{\eta\eta}^c R_{\eta\eta}^c \chi_\eta + 1/18 \bar{A}(\chi_\eta) \bar{C}(\chi_p, \chi_p, \chi_p; 0) R_{\eta\eta}^c R_p^c \chi_p + \bar{A}(\chi_\eta; \varepsilon) \pi_{16} [1/8 \chi_\eta \\
 & - 1/16 \bar{\chi}_1 R_{\eta\eta}^c - 1/8 R_{\eta\eta}^c \chi_\eta - 1/16 R_{\eta\eta}^c \chi_\eta] + \bar{A}(\chi_1) \bar{A}(\chi_3) [-1/72 R_{\eta\eta}^c R_p^c + 1/36 R_{\eta\eta}^c R_{\eta\eta}^c + 1/144 R_{\eta\eta}^c R_p^c] \\
 & - 4 \bar{A}(\chi_{13}) L_1^2 \chi_{13} - 10 \bar{A}(\chi_{13}) L_2^2 \chi_{13} + 1/8 \bar{A}(\chi_{13})^2 - 1/2 \bar{A}(\chi_{13}) \bar{B}(\chi_1, \chi_3; 0, k) \\
 & + 1/4 \bar{A}(\chi_{13}; \varepsilon) \pi_{16} \chi_{13} + 1/4 \bar{A}(\chi_{14}) \bar{A}(\chi_{34}) + 1/16 \bar{A}(\chi_{16}) \bar{A}(\chi_{36}) - 24 \bar{A}(\chi_4) L_1^2 \chi_4 - 6 \bar{A}(\chi_4) L_2^2 \chi_4 \\
 & + 12 \bar{A}(\chi_4) L_4^2 \chi_4 + 1/12 \bar{A}(\chi_4) \bar{B}(\chi_p, \chi_p; 0) (R_{4\eta}^c)^2 \chi_4 + 1/6 \bar{A}(\chi_4) \bar{B}(\chi_p, \chi_\eta; 0) [R_{\eta\eta}^c R_{\eta\eta}^c \chi_4 - 1/18 R_{\eta\eta}^c \chi_4 \\
 & - 1/24 \bar{A}(\chi_4) \bar{B}(\chi_\eta, \chi_\eta; 0) R_{\eta\eta}^c \chi_4 - 1/6 \bar{A}(\chi_4) \bar{B}(\chi_1, \chi_3; 0) R_{\eta\eta}^c R_{\eta\eta}^c \chi_4 + 3/8 \bar{A}(\chi_4; \varepsilon) \pi_{16} \chi_4 \\
 & - 32 \bar{A}(\chi_{46}) L_1^2 \chi_{46} - 8 \bar{A}(\chi_{46}) L_2^2 \chi_{46} + 16 \bar{A}(\chi_{46}) L_4^2 \chi_{46} + \bar{A}(\chi_{46}) \bar{B}(\chi_p, \chi_p; 0) [1/9 \chi_{46} + 1/12 R_{\eta\eta}^c \chi_p \\
 & + 1/36 R_{\eta\eta}^c \chi_4 + 1/9 R_{\eta\eta}^c \chi_6] + \bar{A}(\chi_{46}) \bar{B}(\chi_p, \chi_\eta; 0) [-1/18 R_{\eta\eta}^c \chi_4 - 1/9 R_{\eta\eta}^c \chi_6 + 1/9 R_{\eta\eta}^c \chi_6 + 1/18 R_{\eta\eta}^c \chi_4] \\
 & - 1/6 \bar{A}(\chi_{46}) \bar{B}(\chi_p, \chi_\eta; 0, k) [R_{\eta\eta}^c - R_{\eta\eta}^c] + 1/9 \bar{A}(\chi_{46}) \bar{B}(\chi_\eta, \chi_\eta; 0) R_{\eta\eta}^c \chi_{46} - \bar{A}(\chi_{46}) \bar{B}(\chi_1, \chi_3; 0) [2/9 \chi_{46} \\
 & + 1/9 R_{\eta\eta}^c \chi_6 + 1/18 R_{\eta\eta}^c \chi_4] - 1/6 \bar{A}(\chi_{46}) \bar{B}(\chi_1, \chi_3; 0, k) R_{\eta\eta}^c + 1/2 \bar{A}(\chi_{46}; \varepsilon) \pi_{16} \chi_{46} \\
 & + \bar{B}(\chi_p, \chi_p; 0) \pi_{16} [1/16 \bar{\chi}_1 R_{\eta\eta}^c + 1/96 R_{\eta\eta}^c \chi_p + 1/32 R_{\eta\eta}^c \chi_q] + 2/3 \bar{B}(\chi_p, \chi_p; 0) L_0^2 R_p^c \chi_p \\
 & + 5/3 \bar{B}(\chi_p, \chi_p; 0) L_3^2 R_p^c \chi_p + \bar{B}(\chi_p, \chi_p; 0) L_4^2 [-2 \bar{\chi}_1 \bar{\chi}_{\eta\eta}^{pp} \chi_p - 4 \bar{\chi}_1 R_{\eta\eta}^c \chi_p + 4 \bar{\chi}_1 R_p^c \chi_p + 3 \bar{\chi}_1 R_p^c] \\
 & + \bar{B}(\chi_p, \chi_p; 0) L_5^2 [-2/3 \bar{\chi}_{\eta\eta}^{pp} \chi_p - 4/3 R_{\eta\eta}^c \chi_p^2 + 4/3 R_p^c \chi_p^2 + 1/2 R_{\eta\eta}^c \chi_p - 1/6 R_p^c \chi_q] \\
 & + \bar{B}(\chi_p, \chi_p; 0) L_6^2 [4 \bar{\chi}_1 \bar{\chi}_{\eta\eta}^{pp} + 8 \bar{\chi}_1 R_{\eta\eta}^c \chi_p - 8 \bar{\chi}_1 R_p^c \chi_p] + 4 \bar{B}(\chi_p, \chi_p; 0) L_7^2 (R_p^c)^2 \\
 & + \bar{B}(\chi_p, \chi_p; 0) L_8^2 [4/3 \bar{\chi}_{\eta\eta}^{pp} + 8/3 R_{\eta\eta}^c \chi_p^2 - 8/3 R_p^c \chi_p^2] + \bar{B}(\chi_p, \chi_p; 0)^2 [-1/18 R_{\eta\eta}^c R_p^c \chi_p + 1/18 R_p^c R_{\eta\eta}^c \chi_p \\
 & + 1/288 (R_p^c)^2] + 1/18 \bar{B}(\chi_p, \chi_p; 0) \bar{B}(\chi_\eta, \chi_\eta; 0) [R_{\eta\eta}^c R_p^c \chi_p - R_{\eta\eta}^c R_p^c \chi_p]
 \end{aligned}$$

plus several more pages

Why so long expressions

- Many different quark and meson masses (χ_{ij})
- Charged propagators: $-i G_{ij}^c(k) = \frac{\epsilon_j}{k^2 - \chi_{ij} + i\epsilon} \quad (i \neq j)$
- Neutral propagators: $G_{ij}^n(k) = G_{ij}^c(k) \delta_{ij} - \frac{1}{n_{\text{sea}}} G_{ij}^q(k)$

Why so long expressions

- Many different quark and meson masses (χ_{ij})
- Charged propagators: $-i G_{ij}^c(k) = \frac{\epsilon_j}{k^2 - \chi_{ij} + i\epsilon} \quad (i \neq j)$
- Neutral propagators: $G_{ij}^n(k) = G_{ij}^c(k) \delta_{ij} - \frac{1}{n_{\text{sea}}} G_{ij}^q(k)$

$$-i G_{ii}^q(k) = \frac{R_i^d}{(k^2 - \chi_i + i\epsilon)^2} + \frac{R_i^c}{k^2 - \chi_i + i\epsilon} + \frac{R_{\eta ii}^\pi}{k^2 - \chi_\pi + i\epsilon} + \frac{R_{\pi ii}^\eta}{k^2 - \chi_\eta + i\epsilon}$$

$$R_{jkl}^i = R_{i456jkl}^z, \quad R_i^d = R_{i456\pi\eta}^z,$$

$$R_i^c = R_{4\pi\eta}^i + R_{5\pi\eta}^i + R_{6\pi\eta}^i - R_{\pi\eta\eta}^i - R_{\pi\pi\eta}^i$$

$$R_{ab}^z = \chi_a - \chi_b, \quad R_{abc}^z = \frac{\chi_a - \chi_b}{\chi_a - \chi_c}, \quad R_{abcd}^z = \frac{(\chi_a - \chi_b)(\chi_a - \chi_c)}{\chi_a - \chi_d}$$

$$R_{abcdefg}^z = \frac{(\chi_a - \chi_b)(\chi_a - \chi_c)(\chi_a - \chi_d)}{(\chi_a - \chi_e)(\chi_a - \chi_f)(\chi_a - \chi_g)}$$

Why so long expressions

- Many different quark and meson masses (χ_{ij})
- Charged propagators: $-i G_{ij}^c(k) = \frac{\epsilon_j}{k^2 - \chi_{ij} + i\epsilon} \quad (i \neq j)$
- Neutral propagators: $G_{ij}^n(k) = G_{ij}^c(k) \delta_{ij} - \frac{1}{n_{\text{sea}}} G_{ij}^q(k)$

$$-i G_{ii}^q(k) = \frac{R_i^d}{(k^2 - \chi_i + i\epsilon)^2} + \frac{R_i^c}{k^2 - \chi_i + i\epsilon} + \frac{R_{\eta ii}^\pi}{k^2 - \chi_\pi + i\epsilon} + \frac{R_{\pi ii}^\eta}{k^2 - \chi_\eta + i\epsilon}$$

$$R_{jkl}^i = R_{i456jkl}^z, \quad R_i^d = R_{i456\pi\eta}^z,$$

$$R_i^c = R_{4\pi\eta}^i + R_{5\pi\eta}^i + R_{6\pi\eta}^i - R_{\pi\eta\eta}^i - R_{\pi\pi\eta}^i$$

$$R_{ab}^z = \chi_a - \chi_b, \quad R_{abc}^z = \frac{\chi_a - \chi_b}{\chi_a - \chi_c}, \quad R_{abcd}^z = \frac{(\chi_a - \chi_b)(\chi_a - \chi_c)}{\chi_a - \chi_d}$$


$$R_{abcdefg}^z = \frac{(\chi_a - \chi_b)(\chi_a - \chi_c)(\chi_a - \chi_d)}{(\chi_a - \chi_e)(\chi_a - \chi_f)(\chi_a - \chi_g)}$$

- Relations \implies order of magnitude smaller


Double poles ?

Think quark lines and add gluons everywhere

Full



Quenched




So no resummation at the quark level:

naively a **double pole**

Same follows from inverting the lowest order kinetic terms

Usual resummation from 1PI



$$G_{ij}^n = G_{ij}^0 + G_{ik}^0 (-i) \Sigma_{kl} G_{lj}^0 + G_{ik}^0 (-i) \Sigma_{kl} G_{lm}^0 (-i) \Sigma_{mn} G_{nj}^0 + \dots$$

$$= G^0 (1 + i \Sigma G^0)^{-1}$$

can be done if Σ and G^0 diagonal: usual resummation



$$\frac{1}{p^2 - m^2} \rightarrow \frac{1}{p^2 - m^2 + \Sigma(p^2)}$$

so works in the charged or off-diagonal or $\bar{q}q'$ sector

PQChPT at Two Loop

Use lowest order mass squared: $\chi_i = 2B_0 m_i = m_M^{2(0)}$

Remember: $\chi_i \approx 0.3 \text{ GeV}^2 \approx (550 \text{ MeV})^2 \sim \text{border ChPT}$



Relative corrections: mass²

decay constant

Neutral Masses

Full resummation done by JB, Danielsson to show how to get at the full propagator from one-particle-irreducible diagrams.

Earlier work: Sharpe-Shoresh-Bernard-Golterman
Double poles remain also after resummation.

Actually useful: residue of double pole allows to get at **all** LECs needed for the neutral masses

The eta mass from PQChPT

Expand around the double pole term

$$G_{ij}^n = \frac{-i\mathcal{Z}\mathcal{D}}{(p^2 - M_{ch}^2)^2} + \dots$$

Measure on the lattice via (Sharpe-Shoresh)

$$R_0(t) \equiv \frac{\langle \pi_{ii}(t, \vec{p} = 0) \pi_{jj}(x = 0) \rangle}{\langle \pi_{ij}(t, \vec{p} = 0) \pi_{ji}(x = 0) \rangle} \quad R_0(t \rightarrow \infty) = \frac{\mathcal{D}t}{2M_{ij}},$$

The eta mass from PQChPT

Expand around the double pole term

$$G_{ij}^n = \frac{-i\mathcal{Z}\mathcal{D}}{(p^2 - M_{ch}^2)^2} + \dots$$

Measure on the lattice via (Sharpe-Shoresh)

$$R_0(t) \equiv \frac{\langle \pi_{ii}(t, \vec{p} = 0) \pi_{jj}(x = 0) \rangle}{\langle \pi_{ij}(t, \vec{p} = 0) \pi_{ji}(x = 0) \rangle} \quad R_0(t \rightarrow \infty) = \frac{\mathcal{D}t}{2M_{ij}},$$

From \mathcal{D} can get L_7^r (Sharpe-Shoresh) and has been extended JB, Danielsson to NNLO as well that all LECs relevant for m_η^2 can be had from \mathcal{D} .

A ChPT two-loop timeline: Prehistory

Review paper on Two-Loops: JB, LU TP 06-16
hep-ph/0604043

Dispersive Calculation of the nonpolynomial part in q^2, s, t, u

- Gasser-Meißner: F_V, F_S : 1991 numerical
- Knecht-Moussallam-Stern-Fuchs: $\pi\pi$: 1995 analytical
- Colangelo-Finkemeier-Urech: F_V, F_S : 1996 analytical

Roughly Eurodaphne I

A ChPT two-loop timeline: One mass

Two-Loop Two-Flavour

- Bellucci-Gasser-Sainio: $\gamma\gamma \rightarrow \pi^0\pi^0$: 1994
- Bürgi: $\gamma\gamma \rightarrow \pi^+\pi^-$, F_π , m_π : 1996
- JB-Colangelo-Ecker-Gasser-Sainio: $\pi\pi$, F_π , m_π : 1996-97
- JB-Colangelo-Talavera: $F_{V\pi}(t)$, $F_{S\pi}$: 1998
- JB-Talavera: $\pi \rightarrow \ell\nu\gamma$: 1997
- Gasser-Ivanov-Sainio: $\gamma\gamma \rightarrow \pi^0\pi^0$, $\gamma\gamma \rightarrow \pi^+\pi^-$: 2005-2006

Roughly Eurodaphne II

A ChPT two-loop timeline: ≥ 3 mass

Two-Loops Three flavours

- $\Pi_{VV\pi}, \Pi_{VV\eta}, \Pi_{VVK}$ Kambor, Golowich; Kambor, Dürr; Amorós, JB, Talavera
- $\Pi_{VV\rho\omega}$ Maltman
- $\Pi_{AA\pi}, \Pi_{AA\eta}, F_\pi, F_\eta, m_\pi, m_\eta$ Kambor, Golowich; Amorós, JB, Talavera
- Π_{SS} Moussallam L_4^r, L_6^r
- $\Pi_{VVK}, \Pi_{AAK}, F_K, m_K$ Amorós, JB, Talavera
- $K_{\ell 4}, \langle \bar{q}q \rangle$ Amorós, JB, Talavera L_1^r, L_2^r, L_3^r
- $F_M, m_M, \langle \bar{q}q \rangle (m_u \neq m_d)$ Amorós, JB, Talavera $L_{5,7,8}^r, m_u/m_d$

Roughly Eurodaphne II

A ChPT two-loop timeline: ≥ 3 mass

Two-Loops Three flavours

● $F_{V\pi}, F_{VK^+}, F_{VK^0}$

Post, Schilcher; JB, Talavera L_9^r

● $K_{\ell 3}$

Post, Schilcher; JB, Talavera V_{us}

● $F_{S\pi}, F_{SK}$ (includes σ -terms)

JB, Dhonte L_4^r, L_6^r

● $K, \pi \rightarrow \ell\nu\gamma$

Geng, Ho, Wu L_{10}^r

● $\pi\pi$

JB, Dhonte, Talavera

● πK

JB, Dhonte, Talavera

Roughly Euridice

A ChPT two-loop timeline: ≥ 3 mass

Two-Loops Three flavours

Coming:

- $\eta \rightarrow 3\pi$: status finite for 2/3 of collaboration
- $K_{\ell 3}$ iso: status: diagrams programmed, more time per day needed
- Finite Volume: being thought about
- Finite Volume Two-Flavour: Colangelo-Dürr-Haefeli

Flavianet

A two-loop timeline: \ggg 3 mass

see first part of talk

General Strategy and some comments

- Find enough inputs from experiment
- C_i^r : Ecker's talk
 - kinematical dependence: agree well with single resonance saturation
 - quark mass+kinematical: if vector dominated, seems to be OK
 - quark mass+kinematical: if scalar dominated: which scalars? (not σ)
 - quark masses: which scalars? unrealistically large estimates
- in p^6 physical or lowest order masses: thresholds in right place requires physical

General Strategy and some comments

Inputs:

$K_{\ell 4}$: $F(0), G(0), \lambda$

$m_{\pi^0}^2, m_{\eta}^2, m_{K^+}^2, m_{K^0}^2$

F_{π^+}

F_{K^+} / F_{π^+}

m_s / \hat{m}

L_4^r, L_6^r


C_i^r from single resonance approximation

E865 BNL

em with Dashen violation

24 (26)

$\hat{m} = (m_u + m_d) / 2$



General Strategy and some comments

	fi t 10	same p^4	fi t B	fi t D
$10^3 L_1^r$	0.43 ± 0.12	0.38	0.44	0.44
$10^3 L_2^r$	0.73 ± 0.12	1.59	0.60	0.69
$10^3 L_3^r$	-2.53 ± 0.37	-2.91	-2.31	-2.33
$10^3 L_4^r$	$\equiv 0$	$\equiv 0$	$\equiv 0.5$	$\equiv 0.2$
$10^3 L_5^r$	0.97 ± 0.11	1.46	0.82	0.88
$10^3 L_6^r$	$\equiv 0$	$\equiv 0$	$\equiv 0.1$	$\equiv 0$
$10^3 L_7^r$	-0.31 ± 0.14	-0.49	-0.26	-0.28
$10^3 L_8^r$	0.60 ± 0.18	1.00	0.50	0.54

- ▣ errors are very correlated
- ▣ $\mu = 770$ MeV; 550 or 1000 within errors
- ▣ varying C_i^r factor 2 about errors
- ▣ $L_4^r, L_6^r \approx -0.3, \dots, 0.6 \cdot 10^{-3}$ OK
- ▣ fi t B: small corrections to pion “sigma” term, fi t scalar radius
- ▣ fi t D: fi t $\pi\pi$ and πK thresholds



General Strategy and some comments

	fit 10	same p^4	fit B	fit D
$2B_0\hat{m}/m_\pi^2$	0.736	0.991	1.129	0.958
$m_\pi^2: p^4, p^6$	0.006, 0.258	0.009, $\equiv 0$	-0.138, 0.009	-0.091, 0.133
$m_K^2: p^4, p^6$	0.007, 0.306	0.075, $\equiv 0$	-0.149, 0.094	-0.096, 0.201
$m_\eta^2: p^4, p^6$	-0.052, 0.318	0.013, $\equiv 0$	-0.197, 0.073	-0.151, 0.197
m_u/m_d	0.45 ± 0.05	0.52	0.52	0.50
F_0 [MeV]	87.7	81.1	70.4	80.4
$\frac{F_K}{F_\pi}: p^4, p^6$	0.169, 0.051	0.22, $\equiv 0$	0.153, 0.067	0.159, 0.061

▣▣▣▣ $m_u = 0$ always very far from the fits

▣▣▣▣ F_0 : pion decay constant in the chiral limit

$\pi\pi$




$$a_0^0 = 0.220 \pm 0.005, a_0^2 = -0.0444 \pm 0.0010$$

Colangelo, Gasser, Leutwyler

$$a_0^0 = 0.159 \quad a_0^2 = -0.0454 \text{ at order } p^2$$

$\pi\pi$ and πK



preferred region: fit D: $10^3 L_4^r \approx 0.2$, $10^3 L_6^r \approx 0.0$

General fitting needs more work and systematic studies

Conclusions

- 3 flavour ChPT at 2 loops
 - many calculations done
 - things seem to work but convergence is fairly slow
 - “kinematical” and “vector” C_i^r seem to be OK
 - L_4^r, L_6^r nonzero but reasonable for large N_c
 - $\eta \rightarrow 3\pi$, isobreaking in $K_{\ell 3}$: parts done
- PQChPT at 2 loops: subject just beginning
- $\eta \rightarrow 3\pi$: isospin breaking part needed to p^6 , compare with dispersive (see $K_{\ell 4}$), status: infinities have canceled
- Need to redo fit with all newer data (but then many people not in this audience still use old GL 1985 value of LECs)