

Blocking Technique for Emulating Very Large Polyelectrolytes

Carsten Peterson,* Ola Sommelius,† and Bo Söderberg‡

Department of Theoretical Physics, University of Lund, Sölvegatan 14A, S-223 62 Lund, Sweden

(Received 30 October 1995)

A new Monte Carlo method for computing thermodynamical properties of very large polyelectrolytes is presented. It is based on a renormalization group relating the original polymer to a smaller system, where, in addition to the naively rescaled forces, a corrective nearest-neighbor interaction originating from the short distance Coulomb cutoff is introduced. The method is derived for low T but is in the unscreened case valid for all T . Large polymers with N monomers are emulated by Monte Carlo calculations on smaller systems, $K = N/Q$. The computational gain of the method is Q^3 and is explored with emphasis on room temperature.

PACS numbers: 61.41.+e, 02.70.Lq

The thermodynamics of polyelectrolytes consisting of linear chains of monomers, with covalent harmonic bonding forces and Coulomb interactions, have been extensively studied with Monte Carlo methods. Recently, high statistics results have emerged for relatively long ($N = 2048$) chains [1]. A reliable measurement of, e.g., the end-to-end distance r_{ee} is very CPU time consuming, even when employing state-of-the-art updating methods [2,3].

In order to circumvent the computational bottleneck, and to facilitate estimates of free energies, a variational scheme was proposed [1,4], relying upon an ansatz in the form of a generic Gaussian distribution. For the unscreened case the variational method overestimates r_{ee} by (5–6)% at room temperature, whereas for the screened case the approximation is less accurate.

In this Letter we develop an alternative Monte Carlo (MC) approach, where the chains are “coarse grained” in the spirit of the renormalization group. This is accomplished by using an energy expression for blocked monomers with appropriately renormalized harmonic and Coulomb terms and an additional corrective nearest-neighbor interaction. Superficially, this is reminiscent of the well-known blob picture of de Gennes [5], but the idea is quite different.

The approach is designed for low T . Indeed, at least in the absence of screening, T_r turns out to be low for large N [6]; thus the approach should give good results precisely where it is needed. Our blocking scheme is derived in two ways, using (i) a real space renormalization philosophy, and (ii) a cutoff continuum formulation.

In terms of dimensionless quantities, the energy of a Coulomb chain is given by [1]

$$E = E_G + E_C = \frac{1}{2} \sum_{i=1}^{N-1} \mathbf{r}_{i,i+1}^2 + \sum_i \sum_{j>i} V(r_{ij}), \quad (1)$$

$$V(r) = \frac{e^{-\kappa r}}{r}, \quad (2)$$

with a Boltzmann distribution $\propto \exp(-E/T)$. Here $\mathbf{r}_{ij} = |\mathbf{x}_i - \mathbf{x}_j|$ with \mathbf{x}_i as the monomer positions, while

κ is the inverse Debye screening length. Assuming a length unit of 6 Å, defining the unscreened $N = 2$ equilibrium distance, the temperature in physical units is defined [1] by $e^2 T / 4\pi\epsilon_0\epsilon_r r_0 k_B$. Assuming a dielectric constant $\epsilon_r = 78.3$, $T_r = 0.837808$ will correspond to room temperature (298 K). Similarly, with κ given by $r_0 \sqrt{2e^2 N_A c_s / \epsilon_r \epsilon_0 k_B}$ (298 K), salt concentrations of $c_s = 0.01M$, $0.1M$, and $1.0M$ correspond to $\kappa = 0.1992$, 0.630 , and 1.992 , respectively [1].

In this paper we limit our discussion to the Debye potential model of Eq. (1) for the Coulomb chain, in order to illustrate our ideas; a similar approach can, of course, be applied to other potentials.

Naively, one would coarse grain the chain into blocks of Q monomers, and represent each of them by a single effective position carrying its entire charge. The remaining position variables can then be integrated out (being Gaussian), leaving a system of $K = N/Q$ effective monomers, each carrying a charge Q . They would be connected by harmonic bonds being a factor Q weaker than the original ones. This would be fine, were it not for the divergent short-distance behavior of the Coulomb interaction. This mainly affects neighboring blocks, where the naive effective interaction underestimates the repulsion.

This shortcoming can be remedied by a corrective interaction term between neighboring effective monomers, and we are lead to consider the following blocked energy:

$$E_B^{(K)} = \frac{1}{2Q} \sum_{i=1}^{K-1} r_{i,i+1}^2 + \frac{3}{2} (N - K)T + Q^2 \sum_i \sum_{j>i} V(r_{ij}) + \sum_{i=1}^K W(r_{i,i+1}), \quad (3)$$

where r_{ij} are distances between blocked monomers. The different terms represent, in order of appearance, (1) the naive effective bond energy, (2) a correction to the bond energy, to account for the bond energy of the $N - K$ eliminated position variables, (3) the naive effective interaction, and (4) a corrective nearest-neighbor interaction.

The correction term is determined as follows. In the low temperature limit the chain is a straight line with a slowly varying nearest-neighbor distance, which we approximate by a constant a . For the blocked system, the corresponding nearest-neighbor distance is then $b = Qa$. In this approximation, the true Coulomb energy is given by

$$U_N = \sum_{l=1}^N (N-l)V(la), \quad (4)$$

while the blocked interaction energy (including the correction term) reads

$$U_K = Q^2 \sum_{l=1}^K (K-l)V(lb) + KW(b). \quad (5)$$

The objective is to choose $W(b)$ such that U_N and U_K become identical. For the Coulomb potential of Eq. (1) one obtains for U_N ,

$$\begin{aligned} U_N &= \sum_{l=1}^N (N-l) \frac{e^{-\kappa la}}{la} \approx \frac{N}{a} \sum_{l=1}^{\infty} \frac{e^{-\kappa la}}{l} \\ &= -\frac{N}{a} \ln(1 - e^{-\kappa a}), \end{aligned} \quad (6)$$

and similarly for U_K

$$U_K \approx -\frac{Q^2 K}{b} \ln(1 - e^{-\kappa b}) + KW(b), \quad (7)$$

where only the leading term in the sums have been included. Equating U_N with U_K for $b = Qa$ yields

$$W(r) = -\frac{Q^2}{r} [\ln(1 - e^{-\kappa r/Q}) - \ln(1 - e^{-\kappa r})]. \quad (8)$$

In the absence of screening ($\kappa = 0$) this becomes

$$W(r) = \frac{Q^2 \ln(Q)}{r} \quad (9)$$

representing an enhancement of the nearest-neighbor Coulomb interaction.

The results above are also transparent in a continuum formulation, where the ultraviolet divergence of the Coulomb interaction is regularized via an N -dependent cutoff, as implied by the discrete nature of the original chain. The W term then corrects for the shift in the cutoff. We illustrate this in the unscreened case only.

A continuum formulation of the chain is obtained by replacing the discrete index i by a continuous parameter $\sigma = i/N$, such that $\mathbf{x}_i = \mathbf{x}(\sigma)$. Then the energy of Eq. (1) (in the unscreened case) transforms into

$$\begin{aligned} E &\approx \frac{1}{2N} \int_0^1 \dot{\mathbf{x}}^2(\sigma) d\sigma \\ &+ N^2 \int_0^1 \int_{\sigma+1/N}^1 \frac{d\sigma d\sigma'}{|\mathbf{x}(\sigma') - \mathbf{x}(\sigma)|}. \end{aligned} \quad (10)$$

Were it not for the short-distance cutoff ($1/N$) in σ , needed to regularize the Coulomb potential, all the N dependence could be collected in a global factor N , by making the rescaling $\mathbf{x}(\sigma) \rightarrow N\mathbf{x}(\sigma)$. This corresponds to a mere rescaling of the temperature, and we would have the naive scaling

$$\mathbf{x}_{Qi}^{(QN,QT)} \Leftrightarrow Q\mathbf{x}_i^{(N,T)}. \quad (11)$$

The N -dependent cutoff introduces a scale breaking, and for $K < N$ the last term of Eq. (10) can be rewritten as

$$N^2 \int_0^1 \int_{\sigma+1/N}^1 \frac{d\sigma d\sigma'}{|\mathbf{x}(\sigma') - \mathbf{x}(\sigma)|} = N^2 \int_0^1 \int_{\sigma+1/K}^1 \frac{d\sigma d\sigma'}{|\mathbf{x}(\sigma') - \mathbf{x}(\sigma)|} + N^2 \int_0^1 \int_{\sigma+1/N}^{\sigma+1/K} \frac{d\sigma d\sigma'}{|\mathbf{x}(\sigma') - \mathbf{x}(\sigma)|}. \quad (12)$$

In the low-temperature regime, where the chain is more or less linear, we can use the approximation

$$\mathbf{x}(\sigma') - \mathbf{x}(\sigma) \approx (\sigma' - \sigma)\dot{\mathbf{x}}(\sigma) \quad (13)$$

and obtain for the last term of Eq. (12)

$$\begin{aligned} N^2 \int_0^1 \int_{\sigma+1/N}^{\sigma+1/K} \frac{d\sigma d\sigma'}{|\mathbf{x}(\sigma') - \mathbf{x}(\sigma)|} &\approx N^2 \int_0^1 \int_{\sigma+1/N}^{\sigma+1/K} \frac{d\sigma d\sigma'}{(\sigma' - \sigma)|\dot{\mathbf{x}}(\sigma)|} = N^2 \int_0^1 \frac{d\sigma}{|\dot{\mathbf{x}}(\sigma)|} \int_{1/N}^{1/K} \frac{d\sigma'}{\sigma'} \\ &= N^2 \ln(N/K) \int_0^1 \frac{d\sigma}{|\dot{\mathbf{x}}(\sigma)|}. \end{aligned} \quad (14)$$

Switching back to the discrete representation, the integral becomes the sum of nearest-neighbor interactions; after rescaling with K^2 , it corresponds precisely to the last term of Eq. (3).

The correction to the naive blocked energy function was derived for low temperatures, and the method, as will be demonstrated below, indeed gives excellent results at low T , both for r_{ee} and E_C , for unscreened as well as screened Coulomb interactions.

At higher temperatures the polymer will be less linear, and it is not *a priori* certain that the method will work there. For the *unscreened* case in the high T limit the exact results for r_{ee} and E_C for large N can be expanded in a series in

TABLE I. $r_{ee}(K)/r_{ee}(N)$ as a function of N and $Q = N/K$ at $T = T_r$ for unscreened chains.

N	$Q = 8$	$Q = 4$	$Q = 2$	$Q = 4/3$
40	1.10	1.04	1.01	1.00
80	1.06	1.03	1.01	1.00
160	1.05	1.02	1.00	1.00

$1/T$, yielding [6]

$$\begin{aligned} \langle r_{ee}^2 \rangle &\approx 3(N-1)T + \frac{4}{15} \sqrt{\frac{2}{\pi}} \frac{N^{5/2}}{T^{1/2}}, \\ \langle E_C \rangle &\approx \frac{4}{3} \sqrt{\frac{2}{\pi}} \frac{N^{3/2}}{T^{1/2}}, \end{aligned} \quad (15)$$

while E_G can be determined from the virial identity [1] $2E_G = E_C + 3(N-1)T$. We note that the expressions are consistent with the naive scaling behavior of Eq. (11), in that both r_{ee}/N and E_C/N are functions of T/N only for large N . The relevant small parameter of the high T expansion is obviously N/T . For the blocked system of K effective monomers, the corresponding results are identical to the order shown. Thus in the unscreened case the high T performance is under control.

In the presence of screening, the situation is somewhat different. The correct and blocked high T expansions for E_C differ, and it is actually the contribution from the correction term that dominates [6]. Hence in the screened case the blocking method is reliable only at low T .

We have made extensive numerical evaluations of the blocking approach. All MC runs, both standard and blocked, were performed using the pivot algorithm [2]. Most results are based on 10^4 thermalization sweeps and 10^5 measured configurations. Some results were taken from Ref. [1]. Initially, exploratory comparisons were made in order to find Q , T , and κ ranges where the blocked approach is trustworthy. It turns out that realistic




FIG. 1. r_{ee}/N as a function of N for an unscreened Coulomb chain. Full and dashed lines denote $K = 20$ and $K = 50$ blocked MC runs including logarithmic correction, respectively. Also shown as dotted lines are the naive blocked results for $K = 20$. The squares denote results from original MC runs with N degrees of freedom.




FIG. 2. r_{ee}/N as a function of N for a $\kappa = 0.630$ screened Coulomb chain. Same notation as in Fig. 1.

values for T (T_r) for unscreened chains are within the domain of application even with very high Q values. As expected, in the presence of screening the blocking approach breaks down when the screening length becomes smaller than the blocked resolution. The results are nevertheless encouraging since even with strong screening ($\kappa = 1.992$) $Q \leq 10$ is feasible.

When investigating the power of the method, the relevant parameter is $Q = N/K$. However, in addition, one expects finite- N corrections due to end-point effects. This is illustrated in Table I, where the ratios between r_{ee} computed with blocked and standard method, $r_{ee}(K)$ and $r_{ee}(N)$, respectively, are shown for $T = T_r$ and $\kappa = 0.0$. As can be seen from Table I finite N effects diminish as N increases and should not pose problems for large system sizes, 1000–10000 monomers, at which the method is aimed.

We next compare the power of our blocking approach on r_{ee} and E_C for $T = T_r$ and $\kappa = 0.0$ and 0.630 , respectively. The results for r_{ee} can be found in Figs. 1 and 2. The corresponding energies, which very well approximate the “true” MC values, are found in Table II.

As can be seen from Fig. 1 and Table II the r_{ee} and E_C are amazingly well described by the blocked approach for the unscreened case. The value of r_{ee} is within 10% from the true value even for $Q \approx 100$. For even larger N the situation should be even better with diminishing finite N effects. Indeed, using the blocked method we have estimated r_{ee} and E_C for a $N = 10000$ unscreened chain

TABLE II. Average internal Coulombic energies per monomer for unscreened and screened Coulomb potentials. N and K stand for full MC and blocked with $K = 50$, respectively.

	N	80	160	320	512	1024	2048
$\kappa = 0.0$	$E_C(K)$	2.46	2.76	3.04	3.22	3.48	3.72
	(N)	2.46	2.76	3.04	3.22	3.48	3.73
$\kappa = 0.63$	$E_C(K)$	0.47	0.50	0.54	0.56	0.58	
	(N)	0.47	0.48	0.48	0.48	0.48	

TABLE III. Comparison of $r_{ee}(K, N)$ and $r_{ee}(N, N)$ for $N = 512$ screened chains with $Q = 4$ and 8.

κ	$r_{ee}(N)$	Q	$r_{ee}(K)$	% diff.	Q	$r_{ee}(K)$	% diff.
0.1992	185	4	186	0.6	8	187	1.1
0.630	104	4	105	1.0	8	110	5.8
1.992	61	4	62	1.6	8	68	11.4

with $K = 250$ and find $r_{ee}/N = 2.04$ and $E_C/N = 4.25$. The relative errors should be less than those found for $K = 50$ and $N = 2048$: 3.6% for r_{ee} and 0.3% for E_C . The computational speedup here is substantial, growing as Q^3 .

For the screened case (see Fig. 2), the approximation is less accurate but still good enough to be of practical relevance. In Table III detailed comparisons are made for $Q = 4$ and 8 for different screening strengths. As can be seen there, the blocking gives a reasonable approximation even for $Q = 8$, giving a speedup factor of 512.

The approximation is, of course, not relevant for quantities more local than the blocking allows for.

In summary, we have developed an efficient Monte Carlo blocking scheme that allows for estimating certain thermodynamical quantities of polyelectrolytes with very

substantial computational speedup factors for temperatures around T_r . The method is excellent for unscreened chains, but even with screening non-negligible savings result. Our method is, of course, not confined to Monte Carlo implementations. Other schemes, like molecular dynamics, will benefit from it as well.

We thank Bo Jönsson for valuable comments on the manuscript.

*Electronic address: carsten@thep.lu.se

†Electronic address: ola@thep.lu.se

‡Electronic address: bs@thep.lu.se

- [1] B. Jönsson, C. Peterson, and B. Söderberg, *J. Phys. Chem.* **99**, 1251 (1995).
- [2] M. Lal, *Mol. Phys.* **17**, 57 (1969); N. Madras and A. D. Sokal, *J. Stat. Phys.* **50**, 109 (1988); **50**, 109 (1988).
- [3] A. Irback, *J. Chem. Phys.* **101**, 1661 (1994).
- [4] B. Jönsson, C. Peterson, and B. Söderberg, *Phys. Rev. Lett.* **71**, 376 (1994).
- [5] P. G. deGennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, Ithaca, 1979).
- [6] C. Peterson, O. Sommelius, and B. Söderberg, Report No. LU TP 95-25 (to be published).